

Extension of Strongly Regular Graphs

Ralucca Gera *

Department of Applied Mathematics
Naval Postgraduate School, Monterey, CA 93943
email: rgera@nps.edu, phone (831) 656-2206, fax (831) 656-2355

and

Jian Shen

Department of Mathematics
Texas State University, San Marcos, TX 78666
email: js48@txstate.edu, phone: (512) 245-3740

Submitted: Sep 30, 2007; Accepted: Feb 2, 2008; Published: Feb 11, 2008

Mathematics Subject Classification: 05C75

Abstract

The Friendship Theorem states that if any two people in a party have exactly one common friend, then there exists a politician who is a friend of everybody. In this paper, we generalize the Friendship Theorem. Let λ be any nonnegative integer and μ be any positive integer. Suppose each pair of friends have exactly λ common friends and each pair of strangers have exactly μ common friends in a party. The corresponding graph is a generalization of strongly regular graphs obtained by relaxing the regularity property on vertex degrees. We prove that either everyone has exactly the same number of friends or there exists a politician who is a friend of everybody. As an immediate consequence, this implies a recent conjecture by Limaye et. al.

Key Words: strongly regular graph, Friendship Theorem

1 Introduction and Motivation

In this paper all graphs $G = (V(G), E(G))$ are simple. The neighborhood of a vertex $v \in V(G)$ is $N(v) = \{u : (u, v) \in E(G)\}$. The *join*, denoted $G_1 \vee G_2$, of two graphs G_1 and G_2 is the graph with vertex set $V(G_1) \cup V(G_2)$ and edge set $E(G_1) \cup E(G_2) \cup \{(u, v) : u \in V(G_1) \text{ and } v \in V(G_2)\}$. Also, the *disjoint union*, denoted by $G_1 + G_2$, of two graphs

*Research supported by the Research Initiation Program Grant at the Naval Postgraduate School.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 11 FEB 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Extension of Strongly Regular Graphics				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Applied Mathematics, Monterey, CA, 93943				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

G_1 and G_2 is the graph obtained from G_1 and G_2 with $V(G_1 + G_2) = V(G_1) \cup V(G_2)$ and $E(G_1 + G_2) = E(G_1) \cup E(G_2)$.

We use the notation $d(u) = |N(u)|$ and $\delta(u, v) = |N(u) \cap N(v)|$ to denote the number of neighbors of u (the degree of u) and the number of common neighbors of u and v , respectively. An n -vertex graph is called *strongly regular*, denoted $SRG(n, r, \lambda, \mu)$, if there exist nonnegative integers r, λ, μ such that for all vertices $u, v \in V(G)$,

$$\delta(u, v) = \begin{cases} r & \text{if } u = v; \\ \lambda & \text{if } u \neq v \text{ and } (u, v) \in E(G); \\ \mu & \text{if } u \neq v \text{ and } (u, v) \notin E(G). \end{cases}$$

Many generalizations of strongly regular graphs have been considered in the literature. For example, the t -tuple regular graphs were introduced by Cameron [2]; the NR-regular graphs (also known as the Gamma-Delta regular graphs) were defined by Godsil and McKay [5]; and the strong graphs were defined by Seidel [7]. Since every strongly regular graph has exactly three eigenvalues, graphs with few eigenvalues are also considered as generalizations of strongly regular graphs. In particular, graphs with three eigenvalues were studied by Van Dam [8]; graphs with three Laplacian eigenvalues were studied by Van Dam and Haemers [10]; and graphs with four eigenvalues were studied by Van Dam [9] and by Van Dam and Spence [11].

In this paper, we yet introduce another extension of strongly regular graphs, from a different direction. In 1966, Erdős, Rényi, and Sós [1, 4] proved the following interesting result, commonly referred to as the Friendship Theorem: If $\delta(u, v) = 1$ for any two distinct vertices u, v in a graph G , then $G = K_1 \vee (mK_2)$, where mK_n denotes the disjoint union of m copies of the complete graph on n vertices. A nice interpretation of the theorem is that if any two people in a party have exactly one common friend, then there exists a politician who is a friend of everybody.

We generalize the Friendship Theorem as follows. Let λ and μ be any nonnegative integers. A graph G is called a (λ, μ) -graph if every pair of adjacent vertices have λ common neighbors, and every pair of non-adjacent vertices have μ common neighbors. Thus, (λ, μ) -graphs are generalizations of strongly regular graphs obtained by relaxing the regularity property on vertex degrees. In particular, the Friendship Theorem asserts that $K_1 \vee (mK_2)$ is the only type of $(1, 1)$ -graphs. Since strongly regular graphs have been studied extensively in the literature, we are interested in studying irregular (λ, μ) -graphs. One might assume that there are many (λ, μ) -graphs that are not regular (and thus not strongly regular). To the contrary, we prove that $K_1 \vee (mK_{\lambda+1})$ is the unique type of connected irregular (λ, μ) -graphs. This extends the Friendship Theorem. As an immediate consequence, our result implies the following recent conjecture by Limaye, Sarvate, Stanica, and Young [6] on strongly bi-regular graphs. An n -vertex graph G is called *strongly bi-regular*, denoted $SBRG(n, r, s, \lambda, \mu)$, if G is not regular and there exist nonnegative integers r, s, λ, μ with $r \neq s$ such that for all vertices u, v ,

$$\delta(u, v) = \begin{cases} r \text{ or } s & \text{if } u = v; \\ \lambda & \text{if } u \neq v \text{ and } (u, v) \in E(G); \\ \mu & \text{if } u \neq v \text{ and } (u, v) \notin E(G). \end{cases}$$

Conjecture 1 (Limaye, Sarvate, Stanica, and Young, 2005). *Let G be a connected $SBRG(n, r, s, \lambda, \mu)$. Then $G = K_1 \vee mK_{\lambda+1}$, where $n = m(\lambda + 1) + 1$.*

2 Characterization of Irregular (λ, μ) -graphs

In this section we prove that $K_1 \vee mK_{\lambda+1}$ is the only type of connected irregular (λ, μ) -graphs. A nice interpretation of the result is that if each pair of friends has exactly λ common friends and each pair of strangers have exactly μ ($\mu \geq 1$) common friends in a party, then either everyone has exactly the same number of friends or there exists a politician who is everybody's friend.

Theorem 1. *Suppose G is an irregular (λ, μ) -graph on n vertices. Then one of the following is true:*

- i) $\mu = 0$ and $G = mK_{\lambda+2} + tK_1$ (disjoint union of m copies of $K_{\lambda+2}$ and t copies of K_1), where $n = m(\lambda + 2) + t$.
- ii) $\mu = 1$ and $G = K_1 \vee (mK_{\lambda+1})$, where $n = m(\lambda + 1) + 1$.

Proof. If $\mu = 0$, then G has no pair of vertices with distance two apart. Thus each component of G is a complete graph. Since G is a (λ, μ) -graph, each complete of G is either $K_{\lambda+2}$ or K_1 ; that is, $G = mK_{\lambda+2} + tK_1$. This proves part i) of the theorem.

Suppose now $\mu \neq 0$. For all distinct vertices u, v , define

$$\epsilon(u, v) = \begin{cases} 1 & \text{if } (u, v) \in E(G); \\ 0 & \text{if } (u, v) \notin E(G). \end{cases}$$

Claim 1: $(\mu - \epsilon(u, v))(d(u) - d(v)) = 0$ for any two distinct vertices u, v .

Proof of Claim 1. Let $\epsilon = \epsilon(u, v)$, $A = N(u) - N(v) - \{v\}$, $B = N(u) \cap N(v)$, and $C = N(v) - N(u) - \{u\}$. Let $E(X, Y)$ be the set of edges with one vertex in X and the other vertex in Y , and $E(x, Y)$ be the set of edges with one vertex being $x \in X$ and the other vertex in Y . Then, for each $b \in B$,

$$\begin{aligned} |E(b, A)| &= \begin{cases} |N(b) \cap N(u)| - |E(b, B)| & \text{if } (u, v) \notin E(G); \\ |N(b) \cap N(u)| - |E(b, B)| - 1 & \text{if } (u, v) \in E(G). \end{cases} \\ &= \lambda - |E(b, B)| - \epsilon \end{aligned}$$

and

$$|E(B, A)| = \sum_{b \in B} |E(b, A)| = \sum_{b \in B} (\lambda - \epsilon) - \sum_{b \in B} |E(b, B)| = (\lambda - \epsilon)|B| - 2|E(B, B)|.$$

Similarly,

$$|E(B, C)| = \sum_{b \in B} (|N(b) \cap N(v)| - |E(b, B)| - \epsilon) = (\lambda - \epsilon)|B| - 2|E(B, B)|.$$

Thus

$$|E(A, B)| = |E(B, A)| = |E(B, C)| = |E(C, B)|. \quad (1)$$

Next we count $|E(A, C)|$ in two different ways. First, for each $a \in A$, we have

$$\begin{aligned} |E(a, C)| &= \begin{cases} |N(a) \cap N(v)| - |E(a, B)| & \text{if } (u, v) \notin E(G); \\ |N(a) \cap N(v)| - |E(a, B)| - 1 & \text{if } (u, v) \in E(G). \end{cases} \\ &= \mu - |E(a, B)| - \epsilon. \end{aligned}$$

Thus

$$|E(A, C)| = \sum_{a \in A} |E(a, C)| = \sum_{a \in A} (\mu - |E(a, B)| - \epsilon) = (\mu - \epsilon)|A| - |E(A, B)|.$$

Similarly,

$$|E(C, A)| = \sum_{c \in C} |E(c, A)| = \sum_{c \in C} (\mu - |E(c, B)| - \epsilon) = (\mu - \epsilon)|C| - |E(C, B)|.$$

By (1),

$$(\mu - \epsilon)|A| = |E(A, C)| + |E(A, B)| = |E(C, A)| + |E(C, B)| = (\mu - \epsilon)|C|$$

and thus

$$(\mu - \epsilon)(d(u) - d(v)) = (\mu - \epsilon)((|A| + |B| + \epsilon) - (|C| + |B| + \epsilon)) = (\mu - \epsilon)(|A| - |C|) = 0.$$

This completes the proof of Claim 1.

Recall that $\mu \neq 0$. Then $\mu = 1$, since otherwise, by Claim 1, G would be regular. We now fix a vertex u and define $V_1 = \{v \in V : d(v) = d(u)\}$ and $V_2 = \{v \in V : d(v) \neq d(u)\}$. Since G is irregular, V_1 and V_2 are nonempty and form a partition for V .

Claim 2: Every vertex in V_1 is adjacent to every vertex in V_2 .

Proof of Claim 2. Suppose otherwise $(x, y) \notin E(G)$ for some $x \in V_1$ and some $y \in V_2$. Then $\epsilon(x, y) = 0$. By Claim 1, $d(x) = d(y)$, contradicting the definition for V_1 and V_2 .

Claim 3: Either $|V_1| = 1$ or $|V_2| = 1$.

Proof of Claim 3. Suppose otherwise $|V_1| \geq 2$ and $|V_2| \geq 2$. Let $i = 1, 2$. If there were a pair of non-adjacent vertices u, v in V_i ($i = 1, 2$), then by Claim 2, $\mu = \delta(u, v) \geq |V_{3-i}| \geq 2$, contradicting $\mu = 1$. Thus there is no pair of non-adjacent vertices in each V_i . By Claim 2, $G = K_n$, contradicting the irregularity of G .

By Claim 3, without loss of generality, let $V_1 = \{u\}$. By Claim 2, u is adjacent to every vertex in V_2 . Thus $\mu = 1$ implies that $G[V_2]$, the subgraph of G induced by V_2 , is a disjoint union of complete graphs K_λ . Therefore $G = K_1 \vee (mK_{\lambda+1})$ with $n = m(\lambda + 1) + 1$. This proves the theorem. \square

Remark 1. Since strongly bi-regular graphs are a special case of irregular (λ, μ) -graphs, Theorem 1 implies Conjecture 1 immediately.

Remark 2. We note that our definition of (λ, μ) -graphs extends the concept of all three types of graphs discussed in [3]. Let \overline{G} be the complement graph of G and $N[u] = N(u) \cup \{u\}$ be the closed neighborhood of a vertex u . It can easily be observed that a graph G with n vertices is *uniformly* $(2, r)$ -regular (that is, $|N(u) \cup N(v)| = r$ for each pair of distinct vertices u, v) iff \overline{G} is an $(n - r - 2, n - r)$ -graph; G is *uniformly cl-nbhd* $(2, r)$ -regular (that is, $|N[u] \cup N[v]| = r$ for each pair of distinct vertices u, v) iff \overline{G} is an $(n - r, n - r)$ -graph; G is *k-friendly* (that is, $|N(u) \cap N(v)| = r$ for each pair of distinct vertices u, v) iff G is a (k, k) -graph. Therefore, Theorem 1 implies [3, Corollaries 1, 2, 3, 4].

Acknowledgement. We thank three referees for many valuable comments leading to the clear presentation of the paper.

References

- [1] M. Aigner, G. Ziegler. Proofs from The Book. *Springer-Verlag, Berlin*, 1999.
- [2] P. Cameron. Strongly regular graphs, Chapter 12 in *Selected Topics in Graph Theory*, pp. 337–360, L. Beineke and R. Wilson (Eds.), *Academic Press, New York*, 1978.
- [3] K. Bragan, D. Hoffman, P. Johnson. Inferring regularity by a neglected theorem of Ryser. *Bull. Inst. Combin. Appl.* **50** (2007), 47–51.
- [4] P. Erdős, A. Rényi, V. Sós. On a problem of graph theory. *Studia Sci. Math.* **1** (1966), 215–235.
- [5] C.D. Godsil, B.D. McKay. Graphs with regular neighborhoods. *Combinatorial mathematics, VII (Proc. Seventh Australian Conf., Univ. Newcastle, Newcastle, 1979)*, pp. 127–140, *Lecture Notes in Math.*, **829**, *Springer, Berlin-New York*, 1980.
- [6] N.B. Limaye, D.G. Sarvate, P. Stanica, P. Young. Regular and strongly regular planar graphs. *J. Combin. Math. Combin. Comput.* **54** (2005), 111–127.
- [7] J.J. Seidel. Strongly regular graphs. *1969 Recent Progress in Combinatorics (Proc. Third Waterloo Conf. on Combinatorics, 1968)*, pp. 185–198, *Academic Press, New York*.
- [8] E.R. van Dam. Nonregular graphs with three eigenvalues. *J. Combin. Theory Ser. B* **73** (1998), 101–118.
- [9] E.R. van Dam. Regular graphs with four eigenvalues. *Linear Algebra Appl.* **226/228** (1995), 139–162.
- [10] E.R. van Dam, W.H. Haemers. Graphs with constant μ and $\overline{\mu}$. *Discrete Math.* **182** (1998), no. 1–3, 293–307.
- [11] E.R. van Dam, E. Spence. Small regular graphs with four eigenvalues. *Discrete Math.* **189** (1998), 233–257.